

Influenza and Pneumococcal Vaccines Clinical Practice Guideline

Influenza Vaccine

The following guideline provides recommendations for members who should receive the Influenza vaccine.

Individuals who should receive the vaccine:

- All people including 65 years of age or older
- People with chronic health conditions i.e. heart disease, COPD

Frequency:

- Yearly doses are recommended

Pneumococcal Vaccine

The following guideline provides recommendations for members who should receive pneumococcal vaccine.

For adults 65 or older who do not have an immunocompromising condition, cerebrospinal fluid leak, or cochlear implant and want to receive PPSV23 only	<ul style="list-style-type: none"> • Administer 1 dose of PPSV23 <ul style="list-style-type: none"> • Anyone who received any doses of PPSV23 before age 65 should receive one final dose of the vaccine at age 65 or older. Administer this last dose at least 5 years after the prior PPSV 23 dose
---	---

For adults 65 years or older who do not have an immunocompromising condition, cerebrospinal fluid leak, or cochlear implant and want to receive PCV13 and PPSV23	<ul style="list-style-type: none"> • Administer one dose of PCV13 first, then give one dose of PPSV23 at least one year later <ul style="list-style-type: none"> • Anyone who received any doses of PPSV23 before age 65 should receive one final dose of the vaccine at age 65 or older. Administer this last dose at least 5 years after the prior PPSV23
--	--

This guideline is based on recommendation of Centers of Disease Control cdc.gov/flu/prevent/vaccinations.htm , and cdc.gov/pneumococcal/vaccination.html Rev: 07/2020